

EVA Testing & Prevention Protocol



The following testing and prevention protocol was developed by the EVA Working Group of the American Horse Council and was published in May 1997.

The AHC recommends the adoption of this protocol by all horse breeders as a part of good breeding practice. The protocol provides a practical and realistic approach that permits the use of carrier stallions or infective semen.

Guidelines for breeding a mare to an Equine Arteritis Virus shedding stallion

At least 30 days prior to breeding, the mare should be tested for serum neutralizing antibodies to equine arteritis virus. A blood sample should be submitted to a veterinary medical diagnostic laboratory approved by the USDA to conduct this serological test. Based on that result, the following procedures are recommended.

Antibody Negative (titer less than 1:4) - Non-pregnant Mare

If the mare is found to be serologically negative, she should be vaccinated as soon as possible with the licensed modified live virus vaccine against EVA*. After vaccination, the mare should be isolated for 21 days to allow her time to develop adequate protective immunity against subsequent exposure to the virus and to prevent the minimal risk of spread of the vaccine virus to any susceptible horses with which she might come into contact.

Twenty-one days following vaccination, the mare may be bred to a shedding stallion. She should not be bred to a shedding stallion during that period.

After being bred for the first time to a shedding stallion, the mare should be isolated for 21 days from any horses on the premises serologically negative for antibodies to the virus. Subsequent breedings do not require an additional period of isolation.

Occasionally a mare may be vaccinated against EVA, but for some reason, is not bred that year to a shedding stallion. If this should happen, the mare should be vaccinated again before being bred to a shedding stallion. No isolation is necessary following re-vaccination.

Antibody Negative (titer less than 1:4) - Pregnant Mares

The current licensed modified live virus vaccine against equine viral arteritis* is not approved for use in pregnant mares.

While a mare that is in good health may be vaccinated following parturition, a mare that has had a complicated foaling, or is otherwise not in good health, should not be vaccinated until she has regained her health. The foal should also be in good health and be at least two weeks old before its dam is vaccinated.

There is minimal risk that suckling foals out of serologically negative mares may be exposed to the vaccine virus when the mare is vaccinated against EVA.

Re-vaccination Mares that will be bred to shedding stallions should receive an annual booster vaccination against EVA 21 days prior to being used for breeding purposes. No isolation is necessary following re-vaccination.

Antibody Positive (titer 1:4 or greater) - All Mares

Mares that test serologically positive for antibodies to equine arteritis virus can be bred to a shedding stallion without the need for prior vaccination against EVA. Antibody positive mares that are bred to a shedding stallion by natural cover should be kept separate from other susceptible horses for 24 hours to avoid possible mechanical transmission of virus from voided semen. It should be emphasized that this applies to mares that are serologically positive as a result of previous natural infection with the virus and for mares that have been vaccinated and subsequently bred to a shedding stallion (for management of a vaccinated mare bred the first time to a shedding stallion, see ANTIBODY NEGATIVE [titer of less than 1:4] - Non-Pregnant Mares above). Any vehicle used to transport such mares immediately following breeding to a shedding stallion should be thoroughly cleaned and disinfected prior to transport of susceptible horses.

Guidelines for Breeding Stallions

Prior to the breeding season (at least 60 days is recommended), the stallion should be blood tested for neutralizing antibodies to equine arteritis virus.

Antibody Negative (titer less than 1:4)

If serologically negative, the stallion should be vaccinated with a licensed modified live vaccine against EVA* and isolated for 30 days after vaccination. An annual booster vaccination against EVA should be given on a regular basis every 12 months but no sooner than 30 days prior to being used for breeding.

Antibody Positive (titer 1:4 or greater)

If the stallion is found serologically positive for serum neutralizing antibodies to equine arteritis virus, without written evidence certifying his negative serological status prior to vaccination, he needs to be tested for presence of the carrier (shedding) state. This can be determined by either one of the following methods:

- attempted isolation of equine arteritis virus from two separate ejaculates collected and submitted by an accredited veterinarian to a laboratory approved by the USDA to conduct this test;

OR

- test breeding the stallion to 2 mares serologically negative for antibodies to equine arteritis virus at least twice on each of two consecutive days (four covers) and the mares checked for the development of serum antibodies to the virus 28 days after breeding.

Antibody Positive Non-shedding Stallions

Serologically positive stallions with written certification of negative antibody status prior to vaccination against EVA by a USDA approved laboratory need not be tested for virus shedding.

Stallions serologically positive for antibodies to equine arteritis virus from natural exposure that have previously been tested and found to be non-shedders (non-carriers) of the virus should have written confirmation of their non-shedder status and receive an annual booster vaccination.

EVA Antibody Shedding Positive Stallions

Shedding stallions can be used for commercial breeding provided they are managed in accordance with the above guidelines.

Stallion owners and stallion managers should disclose the shedding status of their stallions to mare owners, breed associations and, where required, to state authorities.

Shedding stallions can be safely bred to adequately immunized mares or to mares that have tested serologically positive for neutralizing antibodies to equine arteritis virus.

Occasionally, shedding stallions will spontaneously stop shedding equine arteritis virus. Owners may wish to retest the semen of shedding stallions from time to time to determine if the stallion is still shedding virus.

Other Recommendations

Teaser Stallions

Teaser stallions should be vaccinated against EVA on an annual basis in accordance with this protocol.

Identification of Carrier (Shedding) Stallions

It is recommended that breed associations publicly disclose the names of those stallions registered with their breed association that are confirmed shedders of equine arteritis virus.

Prevention of the Carrier State

Breeding stallions that are found serologically negative for antibodies to equine arteritis virus should be vaccinated against EVA to prevent the development of the carrier state.

In order to prevent the carrier (shedding) state, especially in those breeds in which the infection is widely prevalent, as well as to prevent equine arteritis virus infection, colts under 270 days of age that are serologically negative for antibodies to equine arteritis virus should be vaccinated against EVA. Written certification of their negative serological status to equine arteritis virus should be obtained before vaccination.

Use of Modified Live Vaccine Against EVA*

It is essential to have written official certification of a horse's negative serological status to equine arteritis virus prior to initial vaccination against this disease.

Stallions and mares that will be bred to shedding stallions should receive an annual booster vaccination against equine arteritis virus prior to being used for breeding purposes.

* ARVAC®, Ft. Dodge Laboratories, Ft. Dodge, Iowa



OIE Reference Center for Equine Viral Arteritis

UNIVERSITY OF KENTUCKY
Department of Veterinary Science
108 Gluck Equine Research Center
Lexington, KY 40546-0099

Ph. (859) 257-4757 Fax (859) 257-8542

Accession # _____
Received _____
Results entered by _____

Contact Information:

Dr. P. Timoney, ext. 81085
Dr. U. Balasuriya, ext. 81124
Ms. K Shuck, ext. 81170

EVA SUBMISSION FORM

(Please print all information)

Veterinarian _____

Owner _____

Address _____

Address _____

Phone (____) _____

Phone (____) _____

FAX (____) _____

Location of Horse _____

Please check if requesting faxed results (\$5 fee)

Email: _____

Signature of Submitting Veterinarian _____

All requested information must be provided to ensure receiving laboratory results

Animal Information

Animal's Registered Name _____ Breed _____

Stallion__ Mare__ Colt__ Filly__ Gelding__ (check one) DOB _____ Tattoo/ID # _____

Vaccinated against EVA? Yes _____ Date(s) _____ No _____

Specimen(s) Submitted

Specimens		Basis for Test
Type of Specimen	Date Collected	
Serum _____	_____	Diagnostic Purposes _____
Semen _____	_____	Export: Animal _____ or Semen _____
Blood EDTA _____	_____	Country _____
Swabs _____	_____	Pre-Vaccination Screen _____
Fetus _____	_____	Breed Registration _____
Age/Dam/Sire _____	_____	Routine Screening _____
Placenta _____	_____	Pre-Purchase Screen _____
Other _____	_____	Surveillance _____
Specify _____	_____	

History (clinical signs, duration, etc., if applicable) _____

LABORATORY RESULTS

For Lab Use Only: Condition _____

Results: Serum _____ (titer _____) Semen: VI _____ PCR _____

Other (Specify) _____

Comments _____

Date Reported _____ Signature _____



Please send copy of results to:
KWPN of North America, KWPN-NA
609 E. Central ❖ P.O. Box O
Sutherlin, OR 97479

Phone: 541-459-3232 ❖ Fax: 541-459-296 7 ❖ E-mail: office@kwpn-na.org



